

Charmed Baryonium

G Cotugno[†], R Faccini^{†,¶}, AD Polosa[¶], C Sabelli^{†,¶}

[¶]INFN Roma, Piazzale A. Moro 2, Roma, I-00185, Italy

[†]Department of Physics, Università di Roma, 'La Sapienza', Piazzale A. Moro 2, Roma, I-00185, Italy

We re-analyze the published data on the $Y(4630) \rightarrow \Lambda_c \bar{\Lambda}_c$ and the $Y(4660) \rightarrow \psi(2S)\pi\pi$ with a consistent Ansatz and we find that the two observations are likely to be due to the *same state* Y_B with $M_{Y_B} = 4660.7 \pm 8.7$ MeV and $\Gamma_{Y_B} = 61 \pm 23$ MeV. Under this hypothesis and reanalyzing also the $e^+e^- \rightarrow J/\psi\pi\pi\gamma_{\text{ISR}}$ spectrum we extract $\mathcal{B}(Y_B \rightarrow \Lambda_c \bar{\Lambda}_c)/\mathcal{B}(Y_B \rightarrow \psi(2S)\pi\pi) = 25 \pm 7$, $\mathcal{B}(Y_B \rightarrow J/\psi\pi\pi)/\mathcal{B}(Y_B \rightarrow \psi(2S)\pi\pi) < 0.46$ @ 90% C.L., $\mathcal{B}(Y(4350) \rightarrow J/\psi\pi\pi)/\mathcal{B}(Y(4350) \rightarrow \psi(2S)\pi\pi) < 3.4 \times 10^{-3}$ @ 90% C.L. and $\mathcal{B}(Y_B \rightarrow \psi(2S)\sigma)/\mathcal{B}(Y_B \rightarrow \psi(2S)f_0) = 2.0 \pm 0.3$. These conclusions strongly support the hypothesis of Y_B being the first observation of a charmed baryonium constituted by four quarks. From the analysis of the mass spectrum we show that $Y(4350)$ and Y_B are respectively consistent with the ground state and first radial excitation of the $\ell = 1$ state.

PACS: 12.39.-x, 12.39.Mk, 13.75.-n

Introduction. The search of the so called baryonia, exotic states originally thought to appear in nucleon-antinucleon systems, has a rather long and sometimes controversial history. Early work in this field can be found in [1] and quite recently there have been new interesting experimental indications as discussed *e.g.* in [2]. In this letter we focus on higher mass scales, namely we refer to the hidden charm sector. We identify as a charmed baryonium a narrow structure, which we call $Y_B(J^{PC} = 1^{--})$, with a decay pattern dominated by the $\Lambda_c \bar{\Lambda}_c$ baryon-antibaryon mode and compare a set of other observables with the expectations of a tetraquark model. Many different interpretations have been proposed in previous works on the nature of the 1^{--} neutral states in the 4 GeV region. The conventional charmonium assignation has been discussed in [3]. Besides the charmonium hybrid interpretation [4] and the *tetraquark* explanation [5], the possibility that $Y(4260)$ could be a threshold effect [6] or a charm mesons molecule [7] have been considered too. Recently [8] and [9] proposed the $Y(4660)$ to be a $D^* \bar{D}_1$ and a $\psi' f_0$ bound state respectively.

Y_B from Belle data. The existence of Y_B stems from our re-analysis of data from the Belle experiment on the decay of the $Y(4660)$ resonance in $\psi(2S)\pi\pi$ [10] and of the $Y(4630)$ in $\Lambda_c \bar{\Lambda}_c$ [11]. We fit the invariant mass spectra in Fig. 1 with a binned likelihood, adopting a consistent signal model: a relativistic Breit-Wigner with comoving width, as detailed in Ref. [12]. Background is parameterized with a second order polynomial multiplied by the phase space.

The individual fits to the charmonium and baryonic modes return $M_{Y_B} = 4661 \pm 9$ MeV, $\Gamma_{Y_B} = 61 \pm 23$ MeV with $\chi^2/\text{d.o.f.} = 7/20$ and $M_{Y_B} = 4661 \pm 14$ MeV, $\Gamma_{Y_B} = 63 \pm 23$ MeV with $\chi^2/\text{d.o.f.} = 51/35$, respectively. The two results are consistent, strongly supporting the hypothesis that the two structures are evidences of the *same resonance*. In the following we will use the results of the first and more accurate, fit as reference. From the same fits we also extract

$$\frac{\mathcal{B}(Y_B \rightarrow \Lambda_c \bar{\Lambda}_c)}{\mathcal{B}(Y_B \rightarrow \psi(2S)\pi^+\pi^-)} = 25 \pm 7, \quad (1)$$

a result which highlights a strong affinity of the Y_B to the baryon-antibaryon decay mode. It is to be noted




FIG. 1: Fit of the cross section from the data of Belle Collaboration [10, 11] for the process $e^+e^- \rightarrow \psi(2S)\pi^+\pi^-$ (left) and $e^+e^- \rightarrow \Lambda_c^+\Lambda_c^-$ (center) and from the BaBar Collaboration [13] for the process $e^+e^- \rightarrow J/\psi\pi^+\pi^-$ (right). The black line represents the fit results, the red one shows the polynomial background.

that omitting interference, the ratio of the peak cross sections is about 11.

Another puzzling aspect of this meson is the absence of the decay mode into $J/\psi\pi^+\pi^-$. In order to try to interpret this observation, we also extract, with the same signal model, an upper limit on the data from BaBar on this decay (see Fig. 1 [13]) and extract, with a bayesian approach, $\mathcal{B}(Y_B \rightarrow J/\psi\pi\pi)/\mathcal{B}(Y_B \rightarrow \psi(2S)\pi\pi) < 0.46$ @ 90% C.L. The presence of the nearby $Y(4260)$ was properly accounted for.

By fitting consistently the $\psi(2S)\pi^+\pi^-$ spectrum in Fig. 1 we also extract the parameters of the $Y(4350)$ resonance. We obtain $M = 4353 \pm 13$ MeV and $\Gamma = 118 \pm 25$ MeV. We also find that $\mathcal{B}(Y(4350) \rightarrow J/\psi\pi\pi)/\mathcal{B}(Y(4350) \rightarrow \psi(2S)\pi\pi) < 3.4 \times 10^{-3}$ @ 90% C.L.

Finally we considered the $\pi\pi$ invariant mass spectrum as measured by Belle in the mass region of the Y_B [10] (see Fig. 2). In absence of background (see Fig. 1), we fit for the sum of two relativistic Breit-Wigner functions describing the σ and $f_0(980)$. The resulting fitted masses and widths are $m_\sigma = (714 \pm 77)$ MeV, $\Gamma_\sigma = (499 \pm 176)$ MeV and $m_{f_0} = (955.3 \pm 4.5)$ MeV, $\Gamma_{f_0} = (14 \pm 15)$ MeV, in agreement with the standard knowledge about the light scalar mesons. Since these mesons are the best currently known candidates for tetraquarks [14], the four-quark nature of the Y_B is naturally suggested. Furthermore we can extract the ratio

$$\mathcal{B}(Y_B \rightarrow \psi(2S)\sigma(600))/\mathcal{B}(Y_B \rightarrow \psi(2S)f_0(980)) = 2.0 \pm 0.3. \quad (2)$$




FIG. 2: Fit of the dipionic mass distribution in the mass region for $\psi(2S)\pi^+\pi^-$: $4.0 \text{ GeV} < m_{\pi^+\pi^-\psi(2S)} < 4.5 \text{ GeV}$ [11] including the contributions of σ and $f_0(980)$. Removing the σ , the lower part of the spectrum gives a sensibly worse fit.

Interpretation of Y_B . The dominance of decay modes with two baryons or with scalar mesons are a strong indication of a $[cq][\bar{c}\bar{q}]$ diquark-antidiquark structure of the Y_B meson. The diagram **A** in Fig. 3 is a representation of a $[cq][\bar{c}\bar{q}]$ tetraquark where the two diquarks are connected by a string neutralizing their color. Both diquarks have positive parity in the ‘single mode configuration’, where the constituent quarks are unexcited with respect to one another. A $J^{PC} = 1^{--}$ state must therefore have odd values of ℓ . The $\ell = 1$ centrifugal potential favors configurations with the diquarks at an higher average relative distance with respect to the $\ell = 0$ case, stretching the color string which eventually breaks as in diagram **B** of Fig. 3 (baryon-antibaryon decay). Breaking the color string in two points rather than in one, as shown in **C** or in **D** in Fig. 3, allows either the




FIG. 3: Phase space allows **A** $\rightarrow$ **B** ($Y \rightarrow \Lambda_c \bar{\Lambda}_c$), **A** $\rightarrow$ **C** ($Y \rightarrow D \bar{D}^*$), **A** $\rightarrow$ **D** ($Y \rightarrow \psi(2S)f_0(980)$)

$D\bar{D}^*$ or the ψf_0 decay. Actually also $\ell = 3$ would be possible assuming the total spin of the diquarks to be $S = 2$ with $S = S_q + S_{\bar{q}}$. Although an analysis of the mass spectrum made along the lines of [17] would exclude this possibility, we will show that a different implementation of the orbital angular momentum dependency of the Hamiltonian would still allow it. We aim at estimating the mass of the Y_B using a non-relativistic constituent quark model combined with a hadron string model. Using the fitted masses of constituent diquarks we will find the spectrum as a function of the orbital angular momentum and radial excitation quantum numbers.

As for the isospin assignment of the tetraquark wavefunction the observed decays seem to favor $I = 0$. On the other hand at the mass scale of the charm quark we could assume that light quark annihilation diagrams are suppressed so that the mass eigenvectors align to the quark mass basis [15]. In the latter case one could have isospin breaking effects as is the case for the $X(3872)$ which turns out to decay in $J/\psi \rho$ and $J/\psi \omega$ at almost the same rate. In a tetraquark interpretation one could explain such decay dynamics making the hypothesis that there are two states [16] of the kind $X_u = [cu][\bar{c}\bar{u}]$ and $X_d = [cd][\bar{c}\bar{d}]$ with a difference in mass of the order of $7 \times \cos(2\theta)$ MeV, θ being the mixing between X_u and X_d ($|\theta| \sim 20^\circ$ in the case of $X(3872)$). From available data we still do not know if this is the case for the Y_B particle although this is not affecting the following considerations.

Angular excitations. The tetraquark picture with two diquarks connected by a color string allows to estimate the mass of the Y_B and its orbital angular momentum excitation level. Considering the generalization of the Chew-Frautschi formula discussed in Selem and Wilczek [18] we find in the mass limit of $M\omega$ ($M = m_1 = m_2$) very large with respect to $\sigma/2\pi$ ¹

$$\mathcal{E} \approx 2M + \frac{3}{(16\pi^2 M)^{1/3}} (\sigma \ell)^{2/3} \quad (3)$$

where $\mathcal{E}$ is the energy of a system made up of two diquarks of mass $M = m_{[cq]}$ held together by a string characterized by a tension σ and spinning with some angular momentum ℓ . The higher ℓ the better this model works as shown in Regge trajectories phenomenology. Taking σ , fitted from Regge slopes of non-charmed hadrons, to be $\sigma \sim 1 \text{ GeV}^2$ and $M = 1933 \text{ MeV}$ [20], we have $\mathcal{E} \approx 4311 \text{ MeV}$ for $\ell = 1$ and $\mathcal{E} \approx 4793 \text{ MeV}$ for $\ell = 3$. This means that if spin-spin interactions are neglected, which is a good approximation at $\ell \neq 0$, and if we consider no radial excitation, the $\ell = 3$ assignment might be favored while the $\ell = 1$ state would better match the $Y(4350)$.

To check the effectiveness of Eq. (3) in estimating the mass of known charmonium resonances having different ℓ it is worth noting that, for $M = 1500 \text{ MeV}$, we get 3500 MeV for the average mass value of the χ_{cJ} system (experimentally located at $\sim 3493 \text{ MeV}$), which has $\ell = 1$, and 3770 MeV for the $\ell = 2$ case consistent with the D -wave nature of the $\psi(3770)$ [21]. The fine structure of the χ_{cJ} system, with levels shifted from the average mass value, is due to spin-orbit and tensor interactions.

Radial excitations. Let us assume that Y_B is a diquark-antidiquark bound state in a Charmoinium-like potential. The difference in mass between the $n_r = 0$ and the first radial excitation $n_r = 1$ can be extracted from the available data on $\chi_{bJ}(2P)$, $\chi_{bJ}(1P)$ states. The mass difference $\Delta M = M_{\chi_b(2P)} - M_{\chi_b(1P)} \simeq 360 \text{ MeV}$ is expected to be roughly the same for charmed states, where instead no data are available for $\ell = 1$ states. Therefore, neglecting spin-orbit terms and spin-spin interactions, the $(n_r = 1, \ell = 1)$ state would be located at 4671 MeV , remarkably close to $Y_B(4660)$, whereas the $(n_r = 0, \ell = 1)$ state would be located at 4311 MeV , close to the $Y(4350)$. We can make an estimate of the mass spectrum also using the constituent quark model discussed in [12]. If we neglect spin-spin and spin-orbit splittings and assume $\ell = 1$ we find $M_Y = 4361 \text{ MeV}$ if no radial excitation is taken into account, and $M_{Y_B} = 4721 \text{ MeV}$ if $n_r = 1$. Therefore we might conclude that $Y_B(4660) = Y(n_r = 1, \ell = 1)$ and $Y(4350) = Y(n_r = 0, \ell = 1)$.

Decays. To refine the model we investigate whether the decay patterns of the $Y(4350)$ and $Y_B(4660)$ are consistent with the assigned quantum numbers: from data we see that both $Y(4350)$ and $Y_B(4660)$ prefer to decay in $\psi(2S)\pi\pi$ rather than in $J/\psi\pi\pi$ even if the latter mode is phase space enhanced. The measured spectra

¹ We perform an infinite mass limit of the the equations of a rotating relativistic string with tension $T = \sigma/2\pi$ and with two masses m_1, m_2 attached at its ends, namely [19]

$$\begin{aligned} \mathcal{E} &= \frac{m_1}{\sqrt{1 - (\omega r_1)^2}} + \frac{m_2}{\sqrt{1 - (\omega r_2)^2}} + \frac{\sigma}{2\pi\omega} \int_0^{\omega r_1} \frac{dv}{\sqrt{1 - v^2}} + \frac{\sigma}{2\pi\omega} \int_0^{\omega r_2} \frac{dv}{\sqrt{1 - v^2}} \\ \ell &= \frac{\omega r_1^2 m_1}{\sqrt{1 - (\omega r_1)^2}} + \frac{\omega r_2^2 m_2}{\sqrt{1 - (\omega r_2)^2}} + \frac{\sigma}{2\pi\omega^2} \int_0^{\omega r_1} \frac{dv v^2}{\sqrt{1 - v^2}} + \frac{\sigma}{2\pi\omega^2} \int_0^{\omega r_2} \frac{dv v^2}{\sqrt{1 - v^2}} \end{aligned}$$

show that the $\pi\pi$ pair comes from phase space (or, equivalently a σ) apart from the case of the $Y_B(4660)$ where there seems to be a 30% component due to $f_0(980)$ (see Eq. 2). We therefore attempt an explanation of the observed Y 's decay pattern describing the S -wave transition $\langle\psi(1S, 2S)\mathbf{a}|Y\rangle$ as

$$\langle\psi(\eta, q)\mathbf{a}(k)|Y(\epsilon, p)\rangle = g \epsilon \cdot \eta \quad (4)$$

where $\mathbf{a} = \sigma, f_0$. We find the eigenfunctions Ψ of the linear part of the Cornell potential $V(r) = -k/r + r/a^2$ ($k = 0.52$, $a = 2.3 \text{ GeV}^{-1}$) for the $\psi(1S, 2S)$ and the charmonium-like Y , namely $\Psi = Y_{\ell, m} R_{n_r, \ell}$. Interpreting the Y 's as charmonium-like bound states made up by a diquark and an antidiquark we can write

$$g \propto \int d^3r R_{(\psi)}^\dagger(r) R_{(Y)}(r) \quad (5)$$

We are interested in estimating the following ratio of decay widths

$$\Gamma_{\mathbf{a}}(Y) \equiv \Gamma(Y \rightarrow \psi(1S)(\pi\pi)_{\mathbf{a}})/\Gamma(Y \rightarrow \psi(2S)(\pi\pi)_{\mathbf{a}}) \quad (6)$$

We find the values reported in the following Table (the square of the matrix element in Eq. (5) weights the three-body phase space where the two pions are the decay products of an intermediate ² scalar resonance; we include only those quantum number combinations which provide mass values close to the experimental ones)

n_r	ℓ	$\Gamma_{\sigma}(Y_B(4660))$	$\Gamma_{f_0}(Y_B(4660))$
1	1	0.07	0.4
0	3	0.2	1.2
n_r	ℓ	$\Gamma_{\sigma}(Y(4350))$	$\Gamma_{f_0}(Y(4350))$
0	1	3.5	—

We report separately the σ and f_0 contributions. The results show that the radial excitation would explain the preference of Y_B tetraquark to decay into $\psi(2S)\sigma$ whereas with this simple model we cannot explain why the $Y(4350)$ seems to prefer the $\psi(2S)$ channel. This aspect remains puzzling.

Conclusions. We reanalyzed the published data on the $Y(4630)$, $Y(4660)$ and $Y(4350)$ and obtained that the first two resonances correspond to a single state, the $Y_B(4660)$. The large baryon decay rate and the decay into $\psi(2S)f_0(980)$ make the Y_B an excellent candidate for a $[cd][\bar{c}\bar{d}]$ diquark-antidiquark bound state suggesting that Y_B represents the first observation of a what we might call a *charmed baryonium*. The fact that the mass of the Y_B is sensibly higher than the baryon-antibaryon threshold excludes any baryon molecule picture. Assuming that the interaction between the two diquarks has the same potential as that between quarks in the charmonium we estimate the mass spectra for the ground state and the first radial excitation for two values of ℓ and find that Y_B best fits the first radial excitation of $\ell = 1$. With the same assumptions we show that the $Y(4350)$ fits the corresponding radial ground state and we show that the absence of the decay into $J/\psi\pi\pi$ of the $Y_B(4660)$ seems compatible with its tetraquark interpretation.

Acknowledgments

We wish to thank Gino Isidori, Antonio Vairo, and Christoph Hanhart for comments and suggestions on the manuscript.

-
- [1] G. C. Rossi and G. Veneziano, Nucl. Phys. B **123**, 507 (1977); L. Montanet, G. C. Rossi and G. Veneziano, Phys. Rept. **63**, 149 (1980).
 - [2] J. P. Dedonder, B. Loiseau, B. El-Bennich and S. Wycech, arXiv:0904.2163 [nucl-th].

² The Breit-Wigner Ansatz we use is a rough approximation for the description of a very broad σ meson whereas is rather suitable for the $f_0(980)$.

- [3] G. J. Ding, J. J. Zhu and M. L. Yan, Phys. Rev. D **77**, 014033 (2008) [arXiv:0708.3712 [hep-ph]].
- [4] S. L. Zhu, Phys. Lett. B **625**, 212 (2005) [arXiv:hep-ph/0507025]; F. E. Close and P. R. Page, Phys. Lett. B **628**, 215 (2005) [arXiv:hep-ph/0507199]; E. Kou and O. Pene, Phys. Lett. B **631**, 164 (2005) [arXiv:hep-ph/0507119].
- [5] L. Maiani, V. Riquer, F. Piccinini and A. D. Polosa, Phys. Rev. D **72**, 031502 (2005) [arXiv:hep-ph/0507062].
- [6] J. L. Rosner, Phys. Rev. D **74**, 076006 (2006) [arXiv:hep-ph/0608102].
- [7] E. Swanson, AIP Conf. Proc. **814**, 203 (2006) [Int. J. Mod. Phys. A **21**, 733 (2006)] [arXiv:hep-ph/0509327]; F. E. Close, *In the Proceedings of 5th Flavor Physics and CP Violation Conference (FPCP 2007), Bled, Slovenia, 12-16 May 2007, pp 020* [arXiv:0706.2709 [hep-ph]].
- [8] F. Close and C. Downum, Phys. Rev. Lett. **102**, 242003 (2009) [arXiv:0905.2687 [hep-ph]]; F. Close, C. Downum and C. E. Thomas, arXiv:1001.2553 [hep-ph].
- [9] F. K. Guo, C. Hanhart and U. G. Meissner, Phys. Lett. B **665**, 26 (2008) [arXiv:0803.1392 [hep-ph]]; F. K. Guo, C. Hanhart and U. G. Meissner, Phys. Rev. Lett. **102**, 242004 (2009) [arXiv:0904.3338 [hep-ph]].
- [10] X. L. Wang *et al.* [Belle Collaboration], "Observation of Two Resonant Structures in e^+e^- to $\pi^+\pi^-\psi(2S)$ via Initial State Radiation at Belle," Phys. Rev. Lett. **99**, 142002 (2007) [arXiv:0707.3699 [hep-ex]].
- [11] G. Pakhlova *et al.* [Belle Collaboration], "Observation of a near-threshold enhancement in the $e^+e^- \rightarrow \Lambda_c^+\Lambda_c^-$ cross section using initial-state radiation," Phys. Rev. Lett. **101**, 172001 (2008) [arXiv:0807.4458 [hep-ex]].
- [12] N. V. Drenska, R. Faccini and A. D. Polosa, Phys. Lett. B **669**, 160 (2008) [arXiv:0807.0593 [hep-ph]].
- [13] B. Aubert *et al.* [BaBar Collaboration], arXiv:0808.1543 [hep-ex].
- [14] L. Maiani, F. Piccinini, A. D. Polosa and V. Riquer, Phys. Rev. Lett. **93**, 212002 (2004) [arXiv:hep-ph/0407017]; G. 't Hooft, G. Isidori, L. Maiani, A. D. Polosa and V. Riquer, Phys. Lett. B **662**, 424 (2008) [arXiv:0801.2288 [hep-ph]].
- [15] G. C. Rossi and G. Veneziano, Phys. Lett. B **70**, 255 (1977); G. C. Rossi and G. Veneziano, Phys. Lett. B **597**, 338 (2004) [arXiv:hep-ph/0404262]; L. Maiani, F. Piccinini, A. D. Polosa and V. Riquer, Phys. Rev. D **70**, 054009 (2004) [arXiv:hep-ph/0407025].
- [16] L. Maiani, A. D. Polosa and V. Riquer, Phys. Rev. Lett. **99**, 182003 (2007) [arXiv:0707.3354 [hep-ph]]; L. Maiani, F. Piccinini, A. D. Polosa and V. Riquer, Phys. Rev. D **71**, 014028 (2005) [arXiv:hep-ph/0412098].
- [17] N. V. Drenska, R. Faccini and A. D. Polosa, Phys. Rev. D **79**, 077502 (2009) [arXiv:0902.2803 [hep-ph]].
- [18] A. Selem and F. Wilczek, arXiv:hep-ph/0602128. See also F. Wilczek, "Diquarks as inspiration and as objects," arXiv:hep-ph/0409168.
- [19] K. Huang, *Quarks Leptons & Gauge Fields*, World Scientific.
- [20] L. Maiani, F. Piccinini, A. D. Polosa and V. Riquer, Phys. Rev. D **71**, 014028 (2005) [arXiv:hep-ph/0412098].
- [21] R. Faccini, "Heavy Quarkonium Spectroscopy," arXiv:0801.2679 [hep-ex].
- [22] B. Aubert *et al.* [BABAR Collaboration], Phys. Rev. Lett. **98**, 212001 (2007) [arXiv:hep-ex/0610057].
- [23] X. L. Wang *et al.* [Belle Collaboration], Phys. Rev. Lett. **99**, 142002 (2007) [arXiv:0707.3699 [hep-ex]].
- [24] L. Maiani, F. Piccinini, A. D. Polosa and V. Riquer, Phys. Rev. D **72**, 031502 (2005) [arXiv:hep-ph/0507062].
- [25] E. Kou and O. Pene, Phys. Lett. B **631**, 164 (2005) [arXiv:hep-ph/0507119].